

HYDROCORTIDERM

Dermatological fluid cream

EXTERNAL USE

Ingredients:

Neomycin sulphate 333.333I.U.
Prednisolone acetate 0.15g
Excipient AD 100g

Mode of Action:

The combination of the different components of Hydrocortiderm provides control of the skin infections of inflammatory or pruriginous origin.

NEOMYCIN: a bacterial and anallergic antibiotic with a broad antibacterial spectrum which is recommended for local use.

PREDNISOLONE: an anti inflammatory corticosteroid which can be used locally in skin infections.

In the form of a dermatologic fluid cream, Hydrocortiderm is easy to apply on the skin, and non greasy to the hands. It is not sticky, irritant or staining.

Indications:

- Local dermatitis
- Complementary treatment of microbial or parasitic dermatitis
- Cutaneous wounds resulting from licking
- 1st degree burns

Dosage – Mode of Administration:

Apply on the lesion 2 to 3 times daily according to the severity of the case, for up to 7 days.

A gentle massage will facilitate the penetration of active principles.

SHAKE BEFORE USE

Presentation:

A white fluid cream containing neomycin and prednisolone which can disperse in water and has a characteristic fragrance.

Plastic bottles of 60gm

Side Effects:

Neomycin may cause some local hypersensitivity which appears as an exudative skin inflammation. Topical prednisolone may exacerbate infection resistant to neomycin, or be responsible for skin atrophy with prolonged use. Discontinue use if this occurs.

Special Precautions:

Wash hands after use. Skin allergy reactions to neomycin and prednisolone have been seen in man after topical use.

Withholding Times:

Meat – 30 days.

Registered pursuant to the ACVM Act 1997,
No A4642

RESTRICTED VETERINARY MEDICINE

Hydrocortiderm

For Animal Treatment Only

